

Author Index

- Abad, C., see Salom, D. 309
 Amiot, J., see Lemieux, L. 399
- Ball, H.L., see Mascagni, P. 375
 Bañó, M.C., see Salom, D. 309
 Bertolini, G., see Mascagni, P. 375
 Beyermann, M., Krause, E. 365
 Bienert, M., Krause, E. 365
 Boppana, V.K.
 — and Miller-Stein, C.
 High-performance liquid chromatographic determination of peptide drugs in biological fluids by means of pre- and post-column fluorescence derivatization techniques 61
 Braco, L., see Salom, D. 309
- Cantini, M., see Libera, L.D. 271
 Carlson, A.D.
 —, Hofer, J.D. and Riggin, R.M.
 Development of an optimized peptide map for recombinant activated human protein C by means of an experimental design strategy 221
 Claessens, H.A., see Štulík, K. 1
 Coeur, C. Le, see Zhao, Q. 201
 Cohen, P., see Rholam, M. 155
 Crimmins, D.L.
 Strong cation-exchange high-performance liquid chromatography as a versatile tool for the characterization and purification of peptides 21
- Desiderio, D.M.
 Analysis of endogenous neuropeptides by reversed-phase high-performance chromatography and mass spectrometry 85
 Deyl, Z.
 —, Mikšík, I., Zicha, J. and Jelínková, D.
 Reversed-phase chromatography of pentosidine-containing CNRr peptides from collagen 257
- Eckert, H., see Koller, M. 31
- Gazdag, M.
 —, Mihályfi, K. and Görög, S.
 Ion-pair RP-HPLC separation of thymopoietin fragments and related peptides 231
 Görög, S., see Gazdag, M. 231
 Guillochon, D., see Lemieux, L. 399
- Hannappel, E., see Huff, T. 239
 Hannappel, E., see Huff, T. 249
 Harada, M.
 High performance liquid chromatographic determination of peptidase activity toward proline-containing peptides 179
 Herraiz, T.
 Sample preparation and reversed phase-high performance liquid chromatography analysis of food-derived peptides 119
 Hofer, J.D., see Carlson, A.D. 221
 Hoffmann, R.
 —, Segal, M. and Otvos, Jr., L.
 Separation of sets of mono- and diphosphorylated peptides by reversed-phase high performance liquid chromatography 327
 Hoffmann, R.
 —, Vasko, M. and Otvos, Jr., L.
 Serum stability of phosphopeptides 319
 Huff, T.
 — and Hannappel, E.
 Oxidation and reduction of thymosin β_4 and its influence on the interaction with G-actin studied by reverse-phase HPLC and post-column derivatization with fluorescamine 249
 Huff, T.
 —, Müller, C. and Hannappel, E.
 HPLC and post-column derivatization with fluorescamine: isolation of actin-sequestering β -thymosins by reversed-phase HPLC 239
 Huisman, T.H.J.
 High performance liquid chromatographic analysis of human hemoglobins and their polypeptide chains: its use in the identification of variants 187
- Irvine, G.B.
 Size-exclusion high-performance liquid chromatography of peptides: a review 387
 Ito, Y., see Ma, Y. 411
- Jelínková, D., see Deyl, Z. 257
- Kai, M.
 —, Ohkura, Y.
 Liquid chromatography of peptides treated with fluorogenic reagents and its application to analyses of opioid peptides, their precursors and related enzymes in rat brain 103
 Kertscher, U., see Mehli, B. 71
 Kline, W., see Schwartz, M. 299

- Kodama, H.
— and Sugahara, K.
Analyses of iminodipeptides containing C-terminal proline or hydroxyproline in biological samples by liquid chromatography/mass spectrometry 141
- Koller, M.
— and Eckert, H.
Derivation of peptides for their determination by chromatographic methods 31
- Krause, E.
—, Rothmund, S., Beyermann, M. and Bienert, M.
Monitoring of α -helical secondary structures in peptides by reversed-phase HPLC of replacement sets 365
- Lemieux, L.
—, Amiot, J., Piot, J.M. and Guillochon, D.
Separation of a casein hydrolyzate by HPSEC with a new mobile phase and characterization of peptides by FABMS 399
- Libera, L.D.
—, Martin, B., Massimino, M.L. and Cantini, M.
Partial purification and biochemical characterization by liquid chromatography of specific myo-mitogenic factor(s) from culture medium of activated macrophages 271
- Ma, Y.
— and Ito, Y.
Recent advances in peptide separation by countercurrent chromatography 411
- Martin, B., see Libera, L.D. 271
- Mascagni, P.
—, Ball, H.L. and Bertolini, G.
Selective purification of synthetic proteins by the use of FMOC- and biotin-based reversible chromatographic probes 375
- Massimino, M.L., see Libera, L.D. 271
- Matuszewski, B., see Schwartz, M. 299
- Mehlis, B.
— and Kertscher, U.
Liquid chromatography/mass spectrometry of peptides of biological samples 71
- Meriluoto, J.
Chromatography of microcystins 277
- Mihályfi, K., see Gazdag, M. 231
- Miksík, I., see Deyl, Z. 257
- Miller-Stein, C., see Boppana, V.K. 61
- Müller, C., see Huff, T. 239
- Ohkura, Y., see Kai, M. 103
- Otvos, Jr., L., see Hoffmann, R. 319
- Otvos, Jr., L., see Hoffmann, R. 327
- Pacáková, V., see Štulík, K. 1
- Péter, A.
— and Tóth, G.
Chromatographic methods for the separation of enantiomers and epimers of β -alkyl amino acids and peptides containing them 335
- Piot, J.M., see Zhao, Q. 201
- Piot, J.M., see Lemieux, L. 399
- Rholam, M.
— and Cohen, P.
Peptides as tools for studying propeptide and proprotein processing 155
- Riggin, R.M., see Carlson, A.D. 221
- Rothmund, S., see Krause, E. 365
- Štulík, K.
—, Pacáková, V., Suchánková, J. and Claessens, H.A.
Stationary phases for peptide analysis by high performance liquid chromatography: a review 1
- Salom, D.
—, Bañó, M.C., Braco, L. and Abad, C.
HPLC in the characterisation of conformational species of linear gramicidins 309
- Schwartz, M.
—, Kline, W. and Matuszewski, B.
Determination of a cyclic hexapeptide (L-743 872), a novel pneumocandin antifungal agent in human plasma and urine by high-performance liquid chromatography with fluorescence detection 299
- Segal, M., see Hoffmann, R. 327
- Suchánková, J., see Štulík, K. 1
- Sugahara, K., see Kodama, H. 141
- Suzuki, T.
—, Watanabe, T. and Toyo'oka, T.
Discrimination of D/L-amino acid in peptide sequence based on fluorescent chiral derivatization by reversed-phase liquid chromatography 357
- Tóth, G., see Péter, A. 335
- Toyo'oka, T., see Suzuki, T. 357
- Vasko, M., see Hoffmann, R. 319
- Watanabe, T., see Suzuki, T. 357
- Zhao, Q.
—, Coeur, C. Le and Piot, J.M.
Analysis of peptides from bovine hemoglobin and tuna myoglobin enzymatic hydrolysate: use of HPLC with on-line second-order derivative spectroscopy for the characterization of biologically active peptides 201
- Zicha, J., see Deyl, Z. 257

